

More Is Not Always Better Backgrounder

CHOOSING WISELY CANADA MISSION

Choosing Wisely Canada is a campaign to help physicians and patients engage in conversations about unnecessary tests, treatments and procedures, and to help physicians and patients make smart and effective choices to ensure high-quality care.

To date, 45 different medical and surgical specialties have joined the Choosing Wisely Canada campaign, representing over 90% of physicians in Canada. Societies representing these specialties have developed evidence-based lists of “top 5” tests and treatments for which there is strong evidence of overuse, waste, or even possible harm to patients. Over 150 physician recommendations have been released across this wide spectrum of specialties. Over 25 corresponding lay language patient materials have also been released to help facilitate the conversation with patients.

For more information on Choosing Wisely Canada, and to view all physician recommendations and patient materials, please visit our website, www.ChoosingWiselyCanada.org. To browse our recommendations and patient materials with ease, download our mobile app by searching for “Choosing Wisely Canada” in your Apple App or Google Play store.

MORE IS NOT ALWAYS BETTER PHILOSOPHY & APPROACH

Since the launch of Choosing Wisely Canada in April 2014, significant effort has gone into educating physicians and mobilizing the health care provider community to adopt the published recommendations. And these efforts are beginning to pay off as impressive results are beginning to emerge across different clinical settings across Canada.

However, these provider-driven efforts are not enough. There is evidence suggesting that patients also play an important role in driving unnecessary care. In fact, in a recent survey of Canadians conducted by Ipsos Reid for Choosing Wisely Canada, 67% of respondents said that “patient demands are responsible for more unnecessary use of health services than are decisions by physicians”. Clearly, the perception that “more is better” needs to be addressed.

The More Is Not Always Better campaign will do the following:

1. Promote the message that in medicine as it is in life, “more is not always better”
2. Educate patients about when they might need a particular test or treatment, and when they don’t
3. Encourage patients to talk with their doctor about unnecessary care.

62% of Canadians agree that there is a significant amount of unnecessary care in the health care system

92% of Canadians believe patients need more support to know which services are really necessary for their health

68% of Canadian family physicians agree that more tools are needed to help them make decisions about which services are inappropriate for their patients

FOUR QUESTIONS TO ASK YOUR DOCTOR

- 1) Do I really need this test, treatment or procedure?
- 2) What are the downsides?
- 3) Are there simpler, safer options?
- 4) What happens if I do nothing?



MORE IS **NOT** ALWAYS BETTER

The same is true for medical tests and treatments.
Talk to your doctor about what you need, and what
you don't. To learn more, visit www.choosingwisely.ca


Choosing
Wisely
Canada



MORE IS NOT ALWAYS BETTER




The same is true for medical tests and treatments.
Talk to your doctor about what you need, and what
you don't. To learn more, visit www.choosingwisely.ca

Choosing
Wisely
Canada 

MORE IS
NOT
ALWAYS
BETTER



The same is true for medical tests and treatments.
Talk to your doctor about what you need, and what
you don't. To learn more, visit www.choosingwisely.ca

Choosing
Wisely
Canada 

Eleven Things Physicians and Patients Should Question

1 Don't do imaging for lower-back pain unless red flags are present.

Red flags include, but are not limited to, severe or progressive neurological deficits or when serious underlying conditions such as osteomyelitis are suspected. Imaging of the lower spine before six weeks does not improve outcomes.

2 Don't use antibiotics for upper respiratory infections that are likely viral in origin, such as influenza-like illness, or self-limiting, such as sinus infections of less than seven days of duration.

Bacterial infections of the respiratory tract, when they do occur, are generally a secondary problem caused by complications from viral infections such as influenza. While it is often difficult to distinguish bacterial from viral sinusitis, nearly all cases are viral. Though cases of bacterial sinusitis can benefit from antibiotics, evidence of such cases does not typically surface until after at least seven days of illness. Not only are antibiotics rarely indicated for upper respiratory illnesses, but some patients experience adverse effects from such medications.

3 Don't order screening chest X-rays and ECGs for asymptomatic or low risk outpatients.

There is little evidence that detection of coronary artery stenosis in asymptomatic patients at low risk for coronary heart disease improves health outcomes. False positive tests are likely to lead to harm through unnecessary invasive procedures, over-treatment and misdiagnosis. Chest X-rays for asymptomatic patients with no specific indications for the imaging have a trivial diagnostic yield, but a significant number of false positive reports. Potential harms of such routine screening exceed the potential benefit.

4 Don't screen women with Pap smears if under 21 years of age or over 69 years of age.

- Don't do screening Pap smears annually in women with previously normal results
- Don't do Pap smears in women who have had a hysterectomy for non-malignant disease

The potential harm from screening women younger than 21 years of age outweighs the benefits and there is little evidence to suggest the necessity of conducting this test annually when previous test results were normal. Women who have had a full hysterectomy for benign disorders no longer require this screening. Screening should stop at age 70 if three previous test results were normal.

5 Don't do annual screening blood tests unless directly indicated by the risk profile of the patient.

There is little evidence to indicate there is value in routine blood tests in asymptomatic patients; instead, this practice is more likely to produce false positive results that may lead to additional unnecessary testing. The decision to perform screening tests, and the selection of which tests to perform, should be done with careful consideration of the patient's age, sex and any possible risk factors.

6 Don't routinely measure Vitamin D in low risk adults.

Because Canada is located above the 35° North latitude, the average Canadian's exposure to sunlight is insufficient to maintain adequate Vitamin D levels, especially during the winter. Therefore, measuring serum 25-hydroxyvitamin D levels is not necessary because routine supplementation with Vitamin D is appropriate for the general population. An exception is made for measuring Vitamin D levels in patients with significant renal or metabolic disease.

7 Don't routinely do screening mammography for average risk women aged 40 - 49.

If, after careful assessment of women less than 50 years of age, their risk profile for breast cancer is low, the benefit of screening mammography is also quite low. Furthermore, for this age group there is a greater risk of a false-positive and consequently undergoing unnecessary or harmful follow-up procedures.

8 Don't do annual physical exams on asymptomatic adults with no significant risk factors.

A periodic physical examination has tremendous benefits; it allows physicians to check on their healthy patients while they remain healthy. However, the benefits of this check-up being done on an annual basis are questionable since many chronic illnesses that benefit from early detection take longer than a year to develop. Preventive health checks should instead be done at time intervals recommended by guidelines, such as those noted by the Canadian Task Force on the Periodic Health Examination.

9 Don't order DEXA (Dual-Energy X-ray Absorptiometry) screening for osteoporosis on low risk patients.

While all patients aged 50 years and older should be evaluated for risk factors for osteoporosis using tools such as the osteoporosis self-assessment screening tool (OST), bone mineral density screening via DEXA is not warranted on women under 65 or men under 70 at low risk.

10 Don't advise non-insulin requiring diabetics to routinely self-monitor blood sugars between office visits.

While self-monitoring of blood glucose (SMBG) for patients with diabetes is recommended by certain groups to help monitor glycemic control, for most adults with type II diabetes who are not using insulin, many studies have shown that routine SMBG does little to control blood sugar over time.

11 Don't order thyroid function tests in asymptomatic patients.

The primary rationale for screening asymptomatic patients is that the resulting treatment results in improved health outcomes when compared with patients who are not screened. There is insufficient evidence available indicating that screening for thyroid diseases will have these results.

How the list was created (1 - 5)

The Canadian Medical Association's (CMA) Forum on General and Family Practice Issues (GP Forum) is a collective of leaders of the General Practice sections of the provincial and territorial medical associations. To establish its *Choosing Wisely Canada* Top 5 recommendations, each GP Forum member consulted with their respective GP Section members to contribute candidate list items. Items from the American Academy of Family Physicians' Choosing Wisely® list were among the candidates. All candidate list items were collated and a literature search was conducted to confirm evidence-based support for the items. GP Forum members discussed which of the thirteen items that resulted should be included. Agreement was found on eight of them. Family physician members of the CMA's e-Panel voted to select five of the eight items. These five items were then approved by the provincial and territorial GP Sections. The College of Family Physicians of Canada is a member observer of the GP Forum and was involved in this list creation process. The first four items on this list are adapted with permission from the Five Things Physicians and Patients Should Question. © 2012 American Academy of Family Physicians.

How the list was created (6 - 11)

The Canadian Medical Association's (CMA) Forum on General and Family Practice Issues (GP Forum) is a collective of leaders of the General Practice sections of the provincial and territorial medical associations. Items 6 - 11 were selected from ten candidate items that were originally proposed for items 1 - 5. GP Forum members discussed which of these items should be included and agreement was found on eight of them. As was done for the first wave, family physician members of the CMA's e-Panel voted to select five of the eight items; however, subsequent discussions by the GP Forum resulted in six items being chosen. Feedback on these six items was then obtained from the provincial/territorial GP Sections. The College of Family Physicians of Canada is a member observer of the GP Forum and was involved in this list creation process.

The GP Forum was dissolved as of August 2015.

Sources

- 1** Canadian Association of Radiologists. The 2012 CAR diagnostic imaging referral guidelines [Internet]. 2012 [cited 2014 Feb 15]. Available from: http://www.car.ca/uploads/standards%20guidelines/car-referralguidelines-c-en_20120918.pdf.
Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: Systematic review and meta-analysis. *Lancet*. 2009 Feb 7;373(9662):463-72.
Ontario Ministry of Health and Long-Term Care (MOHLTC). Excellent care for all - low back pain strategy [Internet]. 2013 [cited 2014 Feb 15]. Available from: http://www.health.gov.on.ca/en/pro/programs/ecfa/action/primary/lb_edutools.aspx.
Physicians of Ontario Collaborating for Knowledge Exchange and Transfer (POCKET). Red and yellow flag indicator cards [Internet]. 2005 [cited 2014 Feb 15]. Available from: <http://www.iwh.on.ca/physicians-network-tool-kit>.
Williams CM, Maher CG, Hancock MJ, McAuley JH, McLachlan AJ, Britt H, et al. Low back pain and best practice care: A survey of general practice physicians. *Arch Intern Med*. 2010 Feb 8;170(3):271-7.
- 2** American Academy of Allergy Asthma and Immunology. Sinus infections account for more antibiotic prescriptions than any other diagnosis [Internet]. 2013 [cited 2014 Feb 15]. Available from: <http://www.aaaai.org/global/latest-research-summaries/Current-JACI-Research/sinus-infection-antibiotic.aspx>.

Desrosiers M, Evans GA, Keith PK, Wright ED, Kaplan A, Bouchard J, et al. Canadian clinical practice guidelines for acute and chronic rhinosinusitis. *Allergy Asthma Clin Immunol.* 2011 Feb 10;7(1):2;1492-7-2.

Hirschmann JV. Antibiotics for common respiratory tract infections in adults. *Arch Intern Med.* 2002 Feb 11;162(3):256-64.

Low D. Reducing antibiotic use in influenza: Challenges and rewards. *Clin Microbiol Infect.* 2008 Apr;14(4):298-306.

Meltzer EO, Hamilos DL. Rhinosinusitis diagnosis and management for the clinician: A synopsis of recent consensus guidelines. *Mayo Clin Proc.* 2011 May;86(5):427-43.

Schumann SA, Hickner J. Patients insist on antibiotics for sinusitis? Here is a good reason to say "no". *J Fam Pract.* 2008 Jul;57(7):464-8.

Smith SR, Montgomery LG, Williams JW Jr. Treatment of mild to moderate sinusitis. *Arch Intern Med.* 2012 Mar 26;172(6):510-3.

- 3** Canadian Association of Radiologists. 2012 CAR diagnostic imaging referral guidelines. Section E: cardiovascular [Internet]. 2012 [cited 2014 Feb 15]. Available from: <http://www.car.ca/uploads/standards%20guidelines/car-referralguidelines-e-en-20121011.pdf>.
Canadian Association of Radiologists. Medical imaging primer with a focus on x-ray usage and safety [Internet]. [Cited 2014 Feb 15]. Available from: http://www.car.ca/uploads/standards%20guidelines/20130128_en_guide_radiation_primer.pdf.
Tigges S, Roberts DL, Vydareny KH, Schulman DA. Routine chest radiography in a primary care setting. *Radiology.* 2004 Nov;233(2):575-8.
U.S. Preventive Services Task Force (USPSTF). Screening for coronary heart disease with electrocardiography [Internet]. 2012 Jul [cited 2014 Feb 15]. Available from: <http://www.uspreventiveservicestaskforce.org/uspstf11/coronarydis/chdfinalrs.htm>.
- 4** Canadian Partnership Against Cancer. Cervical cancer screening guidelines: Environmental scan [Internet]. 2013 Sep [2014 Feb 15]. Available from: http://www.cancerview.ca/idc/groups/public/documents/webcontent/cervical_cancer_enviro_scan.pptx.
Canadian Task Force on Preventive Health Care, Pollock S, Dunfield L, Shane A, Kerner J, Bryant H, et al. Recommendations on screening for cervical cancer. *CMAJ.* 2013 Jan 8;185(1):35-45.
National Institute for Health and Care Excellence. Cervical screening [Internet]. 2010 [cited 2014 Feb 15]. Available from: <http://cks.nice.org.uk/cervical-screening#!scenario recommendation:3>.
- 5** Boland BJ, Wollan PC, Silverstein MD. Yield of laboratory tests for case-finding in the ambulatory general medical examination. *Am J Med.* 1996 Aug;101(2):142-52.
U.S. Preventive Services Task Force. Guide to clinical preventive services: An assessment of the effectiveness of 169 interventions [Internet]. 1989 [cited 2014 Feb 15]. Available from: <http://wonder.cdc.gov/wonder/prevguid/p0000109/p0000109.asp>.
Wians FH. Clinical laboratory tests: Which, why, and what do the results mean?. *Lab Med.* 2009;40:105-13.
- 6** Hanley DA, Cranney A, Jones G, et al. Vitamin D in adult health and disease: a review and guideline statement from Osteoporosis Canada. *CMAJ.* Sep 7 2010;182(12):E610-618.
Toward Optimized Practice (TOP) Working Group for Vitamin D. Guideline for Vitamin D Testing and Supplementation in Adults [Internet]. Edmonton (AB): Toward Optimized Practice; 2012 Oct 31 [cited 2014 Sep 25]. Available from: <http://www.topalbertadoctors.org/download/606/Guideline+for+Vitamin+D+Use+in+Adults+2012+October+31.pdf>.
Guidelines and Protocol Advisory Committee. Vitamin D testing protocol [Internet]. 2013 Jun 1 [cited 2014 Sep 25]. Available from: http://www.bcguidelines.ca/guideline_vitamind.html.
Ontario Association of Medical Laboratories. Guideline for the Appropriate Ordering of Serum Tests for 25-hydroxy Vitamin D and 1,25-dihydroxy Vitamin D [Internet]. 2010 Jun [cited 2014 Sep 25]. Available from: <http://www.oaml.com/PDF/2010/OAML%20Vit%20D%20Guideline%20Jn%20162010%20FINAL.pdf>.
- 7** Canadian Task Force on Preventive Health Care. Screening for breast cancer: Summary of recommendations for clinicians and policymakers [Internet]. 2011 Nov 22 [cited 2014 Sep 25]. Available from: <http://canadiantaskforce.ca/ctfphc-guidelines/2011-breast-cancer/>.
Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement. *Ann. Intern. Med.* Nov 17 2009;151(10):716-726, w-236.
Canadian Task Force on Preventive Health Care. Screening for Breast Cancer. Risk vs. Benefits Poster: For ages 40-49 [Internet]. 2014 [cited 2014 Sep 25]. Available from: <http://canadiantaskforce.ca/perch/resources/breast-cancer-risks-benefits-40-49.pdf>.
Ringash J. Preventive health care, 2001 update: screening mammography among women aged 40-49 years at average risk of breast cancer. *CMAJ.* Feb 20 2001;164(4):469-476.
Tonelli M, Connor Gorber S, Joffres M, et al. Recommendations on screening for breast cancer in average-risk women aged 40-74 years. *CMAJ.* Nov 22 2011;183(17):1991-2001.
- 8** The periodic health examination. Canadian Task Force on the Periodic Health Examination. *Can. Med. Assoc. J.* Nov 3 1979;121(9):1193-1254.
US Preventive Services Task Force Guides to Clinical Preventive Services. The Guide to Clinical Preventive Services 2012: Recommendations of the U.S. Preventive Services Task Force. Rockville (MD): Agency for Healthcare Research and Quality (US); 2012.
Boulware LE, Marinopoulos S, Phillips KA, et al. Systematic review: the value of the periodic health evaluation. *Ann. Intern. Med.* Feb 20 2007;146(4):289-300.
Krogsboll LT, Jorgensen KJ, Gronhoj Larsen C, Gotzsche PC. General health checks in adults for reducing morbidity and mortality from disease: Cochrane systematic review and meta-analysis. *BMJ.* 2012;345:e7191.
Si S, Moss JR, Sullivan TR, Newton SS, Stocks NP. Effectiveness of general practice-based health checks: a systematic review and meta-analysis. *Br. J. Gen. Pract.* Jan 2014;64(618):e47-53.
Blais J, Fournier C, Goulet F, Hanna D, Kossowski A, Laberge C, et al. L'évaluation médicale périodique 2014. Agence de la santé et des services sociaux de Montréal et Collège des médecins du Québec [Internet]. 2014 [cited 2014 Aug 25]. Available from: http://www.cmq.org/fr/MedecinsMembres/Ateliers/~/_media/Files/Guides/EMP-2014.pdf?81425.
- 9** Lim LS, Hoeksema LJ, Sherin K. Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am. J. Prev. Med.* Apr 2009;36(4):366-375.
Papaioannou A, Morin S, Cheung AM, et al. 2010 clinical practice guidelines for the diagnosis and management of osteoporosis in Canada: summary. *CMAJ.* Nov 23 2010;182(17):1864-1873.
Powell H, O'Connor K, Greenberg D. Adherence to the U.S. Preventive Services Task Force 2002 osteoporosis screening guidelines in academic primary care settings. *J Womens Health (Larchmt).* Jan 2012;21(1):50-53.
The International Institute for Clinical Densitometry. 2013 ISCD Official Positions – Adult [Internet]. 2013 [cited 2014 Aug 26]. Available from: <http://www.iscd.org/>.

- 10** Optimal therapy recommendations for the prescribing and use of blood glucose test strips. CADTH Technol Overv. 2010;1(2):e0109.
- Brownlee C. For Diabetics Not on Insulin, Self-Monitoring Blood Sugar Has No Benefit. The Cochrane Library [Internet]. 2012 Jan 19 [cited 2014 Sep 25]. Available from: <http://www.cofah.org/hbns/2012/for-diabetics-not-on-insulin-self-monitoring-blood-sugar-has-no-benefit>.
- Cameron C, Coyle D, Ur E, Klarenbach S. Cost-effectiveness of self-monitoring of blood glucose in patients with type 2 diabetes mellitus managed without insulin. CMAJ. Jan 12 2010;182(1):28-34.
- Gomes T, Juurlink DN, Shah BR, Paterson JM, Mamdani MM. Blood glucose test strips: options to reduce usage. CMAJ. Jan 12 2010;182(1):35-38.
- O'Kane MJ, Bunting B, Copeland M, Coates VE. Efficacy of self monitoring of blood glucose in patients with newly diagnosed type 2 diabetes (ESMON study): randomised controlled trial. BMJ. May 24 2008;336(7654):1174-1177.
- 11** The Canadian Task Force on the Periodic Health Examination. Screening for thyroid disorders and thyroid cancer in asymptomatic adults. The Canadian Guide to Clinical Preventive Health Care [Internet]. 1994;612-18 [cited 2014 Sep 25]. Available from: <http://publications.gc.ca/site/eng/44564/publication.html>
- Screening for thyroid disease: recommendation statement. Ann. Intern. Med. Jan 20 2004;140(2):125-127.
- Surks MI, Ortiz E, Daniels GH, et al. Subclinical thyroid disease: scientific review and guidelines for diagnosis and management. JAMA. Jan 14 2004;291(2):228-238.
- Management of thyroid dysfunction in adults [Internet]. BPJ. 2010 Dec.(22):22-33. [cited 2014 Sep 25]. Available from: <http://www.bpac.org.nz/BPJ/2010/December/thyroid.aspx>.

About The CMA's Forum on General and Family Practice Issues

The Canadian Medical Association's (CMA) Forum on General and Family Practice Issues (GP Forum) is a proud partner of the Choosing Wisely Canada campaign. The GP Forum is a group of family physician leaders in every province and one territory (NWT) in Canada. These individuals are members or chairs of that jurisdiction's Section of General Practice. It also includes representation from the following organizations: the College of Family Physicians of Canada, the Canadian Medical Protective Association, the Society of Rural Physicians of Canada, Canadian Forces Health Services, the Canadian Association of Internes and Residents, the Canadian Federation of Medical Students and one CMA Board member who is a family physician. The primary purpose of the GP Forum is to provide expertise and advice to the CMA on issues concerning primary health care.

Note: The GP Forum was dissolved as of August 2015.

About The College of Family Physicians of Canada

The College of Family Physicians of Canada (CFPC) is a proud partner of the *Choosing Wisely Canada* campaign. The CFPC represents more than 30,000 members across the country. It is the professional organization responsible for establishing standards for the training, certification and lifelong education of family physicians. The College provides quality services and programs, supports family medicine teaching and research, and advocates on behalf of family physicians and the specialty of family medicine. The CFPC accredits postgraduate family medicine training in Canada's 17 medical schools, undergraduate and continuing medical education and encourages the development of research in oncologic surgery. The CSSO believes in facilitating communication between surgeons whose primary interest lies in the field of oncology and encourages the formation of surgical oncology training programs among Canadian Universities.

About Choosing Wisely Canada

Choosing Wisely Canada is a campaign to help physicians and patients engage in conversations about unnecessary tests, treatments and procedures, and to help physicians and patients make smart and effective choices to ensure high-quality care.

For more information on *Choosing Wisely Canada* or to see other lists of Five Things Physicians and Patients Should Question, visit www.choosingwiselycanada.org. Join the conversation on Twitter @ChooseWiselyCA.